

Bracken

Pteridium aquilinum

Plants · Fronds May-Oct · dead litter all winter



Photo by Rasbak, CC BY-SA 3.0, via Wikimedia Commons

A large, coarse fern with tall, triangular, three-times-divided fronds borne singly on stout stalks rather than in a shuttlecock — the way to tell it from every other fern here. Bracken covers extensive areas of hillside, moorland edge, rough grazing and woodland clearing throughout Dumfries and Galloway, turning a distinctive rust-brown in autumn.

BEHAVIOUR

It spreads by deep, tough underground rhizomes, so a whole hillside stand can be one clone, potentially centuries old. It shades out almost everything beneath and its dense litter suppresses regeneration, which makes it a serious problem on grazing and moorland. It is carcinogenic to livestock, and its litter shelters the sheep ticks that carry Lyme disease.

WHEN TO SEE IT

J	F	M	A	M	J	J	A	S	O	N	D
---	---	---	---	---	---	---	---	---	---	---	---

QUICK FACTS

FAMILY	Brackens (Dennstaedtiaceae)
FROND	Up to 2 m or more, borne singly
STATUS	Native; abundant, and invasive on grazing and moorland

Photo by Rasbak, CC BY-SA 3.0, via Wikimedia Commons