

Cinnabar Moth

Tyria jacobaeae

Insects · On the wing May–Aug · flies by day and night



Photo by Charles J. Sharp, CC BY-SA 4.0, via Wikimedia Commons

A striking day- and night-flying moth with dark slate-grey forewings marked by a crimson stripe along the leading edge and two crimson spots, and plain crimson hindwings. Cinnabar moths are common wherever ragwort grows around Dumfries — verges, waste ground, rough pasture and dunes — and are easily disturbed from the vegetation during the day.

BEHAVIOUR

The colouring is a warning: both moth and caterpillar are loaded with toxic alkaloids taken from ragwort, and birds learn quickly to leave them alone. The bright black-and-orange banded caterpillars feed openly and gregariously in mid-summer and can strip a ragwort plant to a bare stem. The moth flies weakly and is often flushed underfoot on a warm afternoon.

WHEN TO SEE IT

J	F	M	A	M	J	J	A	S	O	N	D
---	---	---	---	---	---	---	---	---	---	---	---

QUICK FACTS

FAMILY	Tiger moths (Erebidae, subfamily Arctiinae)
WINGSPAN	3–4.5 cm
STATUS	Native; common wherever ragwort grows

Photo by Charles J. Sharp, CC BY-SA 4.0, via Wikimedia Commons

Nature on your doorstep / Dumfries — part of Love Dumfries · lovedumfries.co.uk/nature